

Advanced Data Modeling

Summer Semester 2008

- Exercises IV -

*To be handed in before 2008-05-26, 23:59 via e-mail to
sschenk@uni-koblenz.de, subject line: [ADM] ...*

1) Instances, Variants and Substitutions

1. Suppose Θ_1 and Θ_2 are substitutions and there exist substitutions σ_1 and σ_2 , such that $\Theta_1 = \Theta_2 \sigma_1$ and $\Theta_2 = \Theta_1 \sigma_2$. Show that there exists a variable-pure substitution γ , such that $\Theta_1 = \Theta_2 \gamma$.
2. Which of the following clauses are Instances or Variants of each other?
 1. $p(x, y, z) :- q(x, y), r(f(z))$
 2. $p(x, b, f(z)) :- q(x, b), r(f(f(z)))$
 3. $p(v, w, f(z)) :- q(v, b), r(f(f(z)))$
 4. $p(z, w, v) :- q(z, w), r(f(v))$
 5. $p(f(x), y, f(z)) :- q(f(x), y), r(f(f(z)))$
 6. $p(f(x), y, z) :- q(f(x), y), r(f(z))$

2) The following Lemma shows that we only need to deal with Herbrand interpretations in order to find a model for any logic program:

Let C be a set of clauses and Σ be any signature containing all symbols used in C . The grounding of C with respect to Σ , denoted C^* is the set of all ground instances of the signature Σ of clauses in C . Let I be a Herbrand interpretation and C be a set of clauses.

Prove that $I \models C$ if and only if $I \models C^*$.

3) Program Completion

1. Let the definition of a predicate symbol p be

$p(y) :- q(y), \text{ not } r(a,y).$
 $p(f(z)) :- \text{ not } q(z).$
 $p(b).$

Give a completion of p .

2. Let P be a normal program and $\text{comp}(P)$ it's completion. Prove that P is a logical consequence of $\text{comp}(P)$. Hint: P is a logical consequence of $\text{comp}(P)$ if $I \models \text{comp}(P) \rightarrow I \models P$